# Lecture 07: Introduction to Normal-Form Games and Nash Equilibria

Lecturer: Anas Barakat October 07, 2025

#### **Abstract**

In this lecture, we provide a succinct introduction to game theory in preparation for our main topic of learning in games in this second part of the course. We define finite normal-form games, pure and mixed Nash equilibria and prove Nash's existence theorem.

Disclaimer: These lecture notes are preliminary notes that will evolve and will be updated over time, this is the first iteration of the course. The treatment is not comprehensive but focuses on some of the main ideas with accompanying proofs.

#### 1 Preamble

Game theory is concerned with analyzing situations involving strategic interactions between several optimizing entities (agents, populations, companies, ...) with different objectives, having an influence on each other via their actions.

Unlike optimization, an agent does not control all the variables that affect her.

An agent's choice of their own controlled decision variable also affects other agents.

In strategic games, we identify three important elements:

- 1. Decision makers: set  $\mathcal{I}$  of entities participating.
  - Terminology: agents, players, users, decision makers.
  - Examples: populations, robots, companies, LLMs, ...
- 2. Choices: Each user  $i \in \mathcal{I}$  is associated to a set of choices.
  - Examples: Machine learning model, routes in a transportation network, ...
- 3. Preferences: Each user  $i \in \mathcal{I}$  is associated with a utility function  $u_i : \mathcal{X} \to \mathbb{R}$  where  $\mathcal{X} := \prod_{i=1}^N \mathcal{X}_i$ .
  - Terminology: utility, preference, payoff, reward, cost, ...
- (System objective: The system of agents might be associated with a performance metric of the form  $W: \mathcal{X} \to \mathbb{R}$ .
  - Examples: social welfare, total cost, aggregated metrics,...)

The interactions between distinct agents depend on multiple considerations:

- The number of agents,
- The incentives of agents: aligned or competing interests,
- Whether agents have the same information about the environment,
- Whether agents must act concurrently or sequentially
- Whether agents can directly communicate with each other.

#### 2 Normal-Form Games

In this section, we introduce normal-form games which is a class of simultaneous single-move games.<sup>1</sup>

#### 2.1 Definition

**Definition 1** (Normal-form games). A strategic game in normal-form is defined by a tuple  $\Gamma := (\mathcal{I}, \{\mathcal{X}_i\}_{i \in \mathcal{I}}, \{u_i\}_{i \in \mathcal{I}})$  where:

- $\mathcal{I}$  is a finite set of players ( $|\mathcal{I}| = n \ge 1$ ),
- $\mathcal{X}_i$  is a non-empty set of strategies for each player  $i \in \mathcal{I}$  ,
- $u_i: \mathcal{X} \to \mathbb{R}$  is a utility function where  $\mathcal{X} := \prod_{i=1}^N \mathcal{X}_i$ .

**Notation.** We use the standard notation  $x = (x_i, x_{-i}) \in \mathcal{X}$  where  $x_{-i} := (x_i)_{i \in \mathcal{I}\{i\}}$ .

**Definition 2.** A game  $\Gamma$  is said to be finite if each strategy set  $\mathcal{X}_i$  is finite.

**Definition 3.** A continuous game is a game  $\Gamma = (\mathcal{I}, \{\mathcal{X}_i\}_{i \in \mathcal{I}}, \{u_i\}_{i \in \mathcal{I}})$  where:

- $\mathcal{I}$  is a finite set of players ( $|\mathcal{I}| = n \ge 1$ ),
- $\mathcal{X}_i$  is a a compact convex subset of a finite dimensional space  $\mathbb{R}^{d_i}$  for each  $i \in \mathcal{I}$ ,
- $u_i: \mathcal{X} \to \mathbb{R}$  (where  $\mathcal{X} := \prod_{i=1}^N \mathcal{X}_i$  is a continuous function mapping an action profile  $x \in \mathcal{X}$  to its associated payoff  $u_i(x) \in \mathbb{R}$ .

We will mostly focus on finite normal-form games.

**Definition 4** (Finite normal-form games). *A finite normal-form game*  $\Gamma = (\mathcal{I}, \{A_i\}_{i \in \mathcal{I}}, \{u_i\}_{i \in \mathcal{I}})$  *is defined by:* 

- a finite set of players  $\mathcal{I} = \{1, \dots, N\}$ ,
- A finite set of actions  $A_i$  for each  $i \in \mathcal{I}$ ,
- A payoff function  $u_i: A \to \mathbb{R}$  for each  $i \in \mathcal{I}$  where  $A := \prod_{i=1}^N A_i$ .

### 2.2 Best response

Each player is seeking to optimize their utility function. Therefore, it is natural to introduce the following best response map which encodes the optimal choice for a player conditioned on the choices of the other players.

**Definition 5** (Best response). The best response of player  $i \in \mathcal{I}$  to the actions  $a_{-i} \in \mathcal{A}_{-i}$  of the other players is a function  $BR^i : \mathcal{A}_{-i} \to 2^{\mathcal{A}_i}$  (also called a correspondence and denoted  $BR^i : \mathcal{A}_{-i} \rightrightarrows \mathcal{A}_i$ ) where

$$BR^{i}(a_{-i}) = \{a_i \in \mathcal{A}_i : u_i(a'_i, a_{-i}) \leq u_i(a_i, a_{-i}), \forall a'_i \in \mathcal{A}_i\}.$$

For any  $\varepsilon \geq 0$ , we define the approximate best response correspondence  $B_{\varepsilon}: \mathcal{A}_{-i} \rightrightarrows \mathcal{A}_{i}$  where

$$BR_{\varepsilon}^{i}(a_{-i}) = \{a_{i} \in \mathcal{A}_{i} : u_{i}(a'_{i}, a_{-i}) \leq u_{i}(a_{i}, a_{-i}) + \varepsilon, \forall a'_{i} \in \mathcal{A}_{i}\}.$$

The best response map is set-valued. We provide an example below for concreteness.

<sup>&</sup>lt;sup>1</sup>Sequential-move games will be discussed later on in the course.

**Example 6.** Consider the following two-player strategic form game with utility functions:

$$\begin{array}{c|ccccc} & L & C & R \\ \hline T & (4,3) & (1,8) & (2,5) \\ M & (2,1) & (0,0) & (-1,1) \\ B & (4,7) & (1,1) & (+1,-1) \\ \end{array}$$

Examples of best response values:  $B_1(L) = \{T, B\}$  and  $B_2(T) = \{C\}$ .

# 3 Nash Equilibria

We are now interested in defining a concept to provide a reasonable description of the collective behavior, as optimality is a priori unclear in a game setting where there are multiple selfish agents with different utility functions. One of the central equilibrium concepts in game theory is the concept of Nash equilibrium. A minimal natural requirement is that no player should be able to improve their payoff (given the behavior of other agents) by switching to a different strategy.

# 3.1 Definition

A Nash equilibrium of a strategic game is a strategy profile such that no player has a unilateral profitable strategy deviation.

**Definition 7** (Nash equilibrium). Let  $\Gamma = (\mathcal{I}, \{A_i\}_{i \in \mathcal{I}}, \{u_i\}_{i \in \mathcal{I}})$  be a finite normal-form game. An action profile  $a^* \in \mathcal{A} = \prod_{i=1}^N \mathcal{A}_i$  is a Nash Equilibrium if:

$$\forall i \in \mathcal{I}, \quad \forall a_i \in \mathcal{A}_i, \quad u_i(a_i, a_{-i}^*) \le u_i(a_i^*, a_{-i}^*), \tag{1}$$

or equivalently:

$$\forall i \in \mathcal{I}, \quad \forall a_i \in \mathcal{A}_i, \quad a_i^* \in BR^i(a_{-i}^*).$$
 (2)

For any  $\varepsilon \geq 0$ , an  $\varepsilon$ -Nash equilibrium is an action profile  $a^* \in \mathcal{A}$  s.t.

$$\forall i \in \mathcal{I}, \quad \forall a_i \in \mathcal{A}_i, \quad u_i(a_i, a_{-i}^*) \le u_i(a_i^*, a_{-i}^*) + \varepsilon, \tag{3}$$

or equivalently:

$$\forall i \in \mathcal{I}, \quad \forall a_i \in \mathcal{A}_i, \quad a_i^* \in BR_{\varepsilon}^i(a_{-i}^*). \tag{4}$$

We provide a few comments regarding this definition and the interpretation of NE:

- Each player  $i \in \mathcal{I}$  is playing a best response to the actions of the other players.
- In a Nash equilibrium, no player regrets their action choice (i.e. could have played a better action given the actions of others).
- Only *relative* preferences matter in the definition of NE, i.e. deviations  $u_i(a_i, a_{-i}^*) u_i(a_i^*, a_{-i}^*)$  for  $a_i \in \mathcal{A}_i$ .

**Remark 8.** The definition extends naturally to continuous games by replacing  $A_i$  by  $X_i$ .

A few natural questions arise:

- Do NE (always) exist?
- When they do exist, are they unique in general? If not, are they equal in terms of payoffs? We provide a few answers by looking at a few classical examples.

## 3.2 Examples and table representation

**Example 9** (Prisoner's dilemma). Two suspects are independently interrogated. They have the choice between confessing to committing the crime (including reporting their partner) or not confessing. If both cooperate, both are sentenced to 1 year jail. If they both defect, they are both sentenced to 3 years. Otherwise, if one cooperates and the other defects, the payoffs are respectively -4 and 0.

$$\begin{array}{c|cccc}
 & C & D \\
\hline
C & (-1,-1) & (-4,0) \\
D & (0,-4) & (-3,-3)
\end{array}$$

It can be verified that (D, D) is the unique Nash equilibrium for this game. Note for this that action D strictly dominates action C for both players. The suspects' most desirable collective behavior is (C, C) but it is not stable in the sense of NE.

**Example 10** (Bach or Stravinsky). A couple would like to go to a music concert to listen either to Bach or Stravinsky. The couple prefers to go together but the partners have different preferences: one prefers Bach whereas the other prefers Stravinsky. A payoff matrix modeling these preferences is given as follows:

$$\begin{array}{c|cccc}
 & B & S \\
\hline
B & (2,1) & (0,0) \\
S & (0,0) & (1,2)
\end{array}$$

Notice that there are two Nash equilibria (B, B) and (S, S).

**Example 11** (Matching Pennies). Each player simultaneously chooses either Heads (H) or Tails (T). If the choices match (both choose H or both choose T), Player 1 wins and Player 2 loses. If the choices differ, Player 2 wins and Player 1 loses.

$$\begin{array}{c|cccc} & H & T \\ \hline H & (1,-1) & (-1,1) \\ T & (-1,1) & (1,-1) \end{array}$$

Note that there does not exist any NE! There is always a profitable unilateral strategy deviation.

This example motivates the use of a randomized strategy.

# 3.3 Mixed extension of normal-form games

The mixed extension of a finite game  $\Gamma = (\mathcal{I}, \{\mathcal{A}_i\}_{i \in \mathcal{I}}, \{\sqcap_i\}_{i \in \mathcal{I}})$  is a compact continuous (multilinear) game  $\tilde{\Gamma}$  (which we will denote  $\Delta(\Gamma)$ ) defined by the tuple  $(\mathcal{I}, \{\mathcal{X}_i\}_{i \in \mathcal{I}}, \{\tilde{u}_i\}_{i \in \mathcal{I}})$  where:

- for all  $i \in \mathcal{I}$ ,  $\mathcal{X}_i = \Delta(\mathcal{A}_i)$  where  $\Delta(\mathcal{A}_i) = \{x_i \in \mathbb{R}^{|\mathcal{A}_i|} : x_{i,a_i} \geq 0, \sum_{a_i \in \mathcal{A}_i} x_{i,a_i} = 1\}$  is the set of probability distributions over the finite set  $\mathcal{A}_i$  with cardinality  $|\mathcal{A}_i|$ ,
- for all  $i \in \mathcal{I}$ , the payoff of player i in  $\tilde{\Gamma}$  is the multilinear extension of the utility function  $u_i$  (in the original game  $\Gamma$ ) which we denote by  $\tilde{u}_i : \mathcal{X} \to \mathbb{R}$  defined for any strategy profile  $x \in \mathcal{X} = \prod_{i=1}^N \Delta(\mathcal{A}_i)$  by:

$$\tilde{u}_i(x) = \sum_{a = (a_1, \dots, a_N) \in \mathcal{A}} \left( \prod_{j=1}^N x_j(a_j) \right) \cdot u_i(a) = \mathbb{E}_{a_1 \sim x_1, \dots, a_n \sim x_n} [u_i(a_1, \dots, a_N)].$$
 (5)

With a slight abuse of notation, we will reuse the notation  $u_i$  for  $\tilde{u}_i$ .

**Notation.** We will use the notation  $u_i(a_1, x_{-i})$  for the payoff of player i when player 1 chooses action  $a_1$  while other players select strategies  $x_{-i}$ .

**Remark 12** (Mixed vs correlated strategies). Note that a mixed strategy as we define it here is a strategy  $x \in \prod_{i=1}^N \Delta(\mathcal{A}_i)$  (product of independent distributions) whereas we can define a correlated strategy as  $x \in \Delta(\prod_{i=1}^N \mathcal{A}_i)$  (distribution over the product space). We will come back to this distinction later on in the next lectures.

**Example 13** (Rock-paper-scissors). For any a, b > 0, define the following game with payoffs:

$$\begin{array}{c|ccccc} & R & P & S \\ \hline R & (0,0) & (a,-b) & (-b,a) \\ P & (-b,a) & (0,0) & (a,-b) \\ S & (a,-b) & (-b,a) & (0,0) \\ \end{array}$$

For the fully-mixed strategy  $x_1 = (\frac{1}{4}, \frac{1}{4}, \frac{1}{2})$  and the strategy  $x_2 = (\frac{1}{2}, \frac{1}{2}, 0)$ , compute the expected payoff  $u_1(x_1, x_2)$ .

**Definition 14** (Mixed Nash equilibrium). *A mixed Nash equilibrium of*  $\Gamma$  *is a Nash equilibrium of the mixed extension*  $\Delta(\Gamma)$ .

In the RPS game (example 13) with a = b = 1, the only Nash equilibrium is the uniform strategy for all players. In particular, there is no NE in pure strategies (pure NE).

Naturally, the definition of the best response correspondence can also be generalized.

**Definition 15** (Best response correspondence). The best response correspondence  $BR : \mathcal{X} \rightrightarrows \mathcal{X}$  which maps to any strategy  $x \in \mathcal{X}$  the subset  $\prod_{i=1}^{N} BR^{i}(x_{-i})$  of  $\mathcal{X}$ , i.e.

$$BR(x) = BR(x_1, \dots, x_N) = (BR^1(x_{-1}), \dots, BR^N(x_{-N})).$$
 (6)

Using this definition we obtain a fixed-point characterization of NE (mixed or pure):

**Proposition 16.** A strategy  $x^*$  is a NE if and only if  $x^* \in BR(x^*)$ .

**Remark 17** (Computational complexity). It can be shown that computation of fixed points of continuous functions and computation of NE are computationally equivalent, i.e. each problem can be reduced to the other in polynomial time.

# 4 Existence of Mixed Nash Equilibria

### 4.1 Nash's theorem

**Theorem 18** (Nash (1951)). Every finite game  $\Gamma$  has (at least) a mixed Nash equilibrium.

#### 4.2 Proof

The original proof (Nash, 1950) is based on Kakutani's fixed point theorem (Kakutani, 1941).

**Theorem 19** ((Kakutani, 1941)). Let C be a non-empty convex and compact subset of a normed vector space and let F be a correspondence from C to C such that:

- 1. For all  $c \in C$ , F(c) is convex, compact, and non-empty;
- 2. The graph  $\Gamma = \{(c,d) \in C \times C : d \in F(c)\}$  of F is closed.

Then,  $\{c \in C : c \in F(c)\}$  is non-empty and compact.

**Remark 20.** In 1952, Glicksberg (1952) and Fan (1952) independently generalized Kakutani's theorem to any Hausdorff locally convex topological vector space.

A year later, Nash (Nash, 1951) proposed an alternative proof based on Brouwer's fixed point theorem. We will present the latter in the following<sup>2</sup>.

**Theorem 21** (Brouwer's fixed point theorem). *Let* C *be a non-empty convex and compact subset of a finite-dimensional Euclidean space. Then any continuous function*  $f: C \to C$  *has a fixed point.* 

First, define for any  $x \in \prod_{i=1}^{N} \Delta(A_i)$  and any  $a_i \in A_i$ ,

$$r_{i,a_i}(x) = u_i(a_i, x_{-i}) - u_i(x)$$
. (7)

We introduce Nash's map  $\varphi: \prod_{i=1}^{N} \Delta(A_i) \to \prod_{i=1}^{N} \Delta(A_i)$  defined for every  $x \in \prod_{i=1}^{N} \Delta(A_i)$  by:

$$\varphi_{i,a_i}(x) := \frac{x_{i,a_i} + [r_{i,a_i}(x)]^+}{1 + \sum_{a_i' \in A_i} \left[ r_{i,a_i'}(x) \right]^+},$$
(8)

for all player  $i \in [n]$  and action  $a_i \in A_i$ . We use the notation  $[r]^+ := \max\{0, r\}$ .

**Proposition 22** (Utility improvement under Nash map). *For any strategy profile*  $x \in \prod_{i=1}^{N} \Delta(A_i)$ , and any player  $i \in [n]$ ,

$$u_{i}\left(\varphi_{i}(x), x_{-i}\right) - u_{i}(x_{i}, x_{-i}) = \frac{\sum_{a_{i} \in \mathcal{A}_{i}} \left(\left[r_{i, a_{i}}(x)\right]^{+}\right)^{2}}{1 + \sum_{a_{i} \in \mathcal{A}_{i}} \left[r_{i, a_{i}}(x)\right]^{+}}.$$
(9)

- Useful proof technique.
- Interpretation: if a player has an incentive to unilaterally deviate, then the Nash map unilaterally increases that player's utility.
- Even if a single action is profitable, then the Nash map unilaterally strictly increases for the utility of that player.

*Proof.* Fix  $i \in \mathcal{I}$  and introduce the following shorthand notations for  $x \in \mathcal{X}$ ,  $a_i \in \mathcal{A}_i$ ,

$$r_{i,a_i} := r_{i,a_i}(x)$$
,  $u_{i,a_i} := u_i(a_i, x_{-i})$ ,  $x'_{i,a_i} := \varphi_{i,a_i}(x_1, \dots, x_n) = \frac{x_{i,a_i} + r^+_{i,a_i}}{1 + \sum_{a'_i \in A_i} r^+_{i,a'_i}}$ . (10)

<sup>&</sup>lt;sup>2</sup>We follow here the exposition in the lecture notes of G. Farina (Farina, 2024)

The utility improvement under Nash's map is computed as follows:

$$\begin{aligned} u_{i}\left(\varphi_{i}(x),x_{-i}\right) - u_{i}(x_{i},x_{-i}) &= u_{i}\left(x'_{i},x_{-i}\right) - u_{i}(x_{i},x_{-i}) \\ &= \sum_{a_{i} \in A_{i}} u_{i,a_{i}} \cdot \left(x'_{i,a_{i}} - x_{i,a_{i}}\right) \\ &= \sum_{a_{i} \in A_{i}} u_{i,a_{i}} \cdot \left(\frac{x_{i,a_{i}} + r^{+}_{i,a_{i}}}{1 + \sum_{a'_{i} \in A_{i}} r^{+}_{i,a'_{i}}} - x_{i,a_{i}}\right) \\ &= \sum_{a_{i} \in A_{i}} u_{i,a_{i}} \cdot \frac{r^{+}_{i,a_{i}} - \sum_{a'_{i} \in A_{i}} r^{+}_{i,a'_{i}}}{1 + \sum_{a'_{i} \in A_{i}} r^{+}_{i,a'_{i}}} \\ &= \frac{1}{1 + \sum_{a'_{i} \in A_{i}} r^{+}_{i,a'_{i}}} \left(\sum_{a_{i} \in A_{i}} r^{+}_{i,a_{i}} \cdot u_{i,a_{i}} - \sum_{a'_{i} \in A_{i}} \left(r^{+}_{i,a'_{i}} \sum_{a_{i} \in A_{i}} x_{i,a_{i}} \cdot u_{i,a_{i}}\right)\right) \\ &= \frac{1}{1 + \sum_{a'_{i} \in A_{i}} r^{+}_{i,a'_{i}}} \left(\sum_{a_{i} \in A_{i}} r^{+}_{i,a_{i}} \cdot \left(u_{i,a_{i}} - \sum_{a'_{i} \in A_{i}} u_{i,a'_{i}} \cdot x_{i,a'_{i}}\right)\right) \\ &= \frac{1}{1 + \sum_{a'_{i} \in A_{i}} r^{+}_{i,a'_{i}}} \left(\sum_{a_{i} \in A_{i}} r^{+}_{i,a_{i}} \cdot r_{i,a_{i}}\right) \\ &= \frac{\sum_{a_{i} \in A_{i}} \left(\left[r_{i,a_{i}}(x)\right]^{+}\right)^{2}}{1 + \sum_{a_{i} \in A_{i}} \left[r_{i,a_{i}}(x)\right]^{+}}, \end{aligned}$$

where the last inequality follows from observing that  $z^+ \cdot z = (z^+)^2$  for all  $z \in \mathbb{R}$ .

**Proposition 23.** A strategy profile  $x \in \mathcal{X}$  is a Nash equilibrium if and only if it is a fixed point of the Nash improvement function  $\varphi$ .

*Proof.* ( $\Longrightarrow$ ) If x is a NE, then by definition for all  $i \in [n]$  and  $a_i \in \mathcal{A}_i$ , we have  $r_{i,a_i}(x) \leq 0$ , hence  $[r_{i,a_i}(x)]^+ = 0$ . Therefore for all  $i \in \mathcal{I}$  and all  $a_i \in \mathcal{A}_i$ ,

$$\varphi_{i,a_i}(x) = \frac{x_{i,a_i} + [r_{i,a_i}(x)]^+}{1 + \sum_{a_i' \in A_i} [r_{i,a_i'}(x)]^+} = x_{i,a_i}.$$
 (12)

 $(\Leftarrow)$  Conversely, suppose that x is a fixed point of  $\varphi$ , i.e.  $x = \varphi(x)$ . Then, for all  $i \in \mathcal{I}$ , it follows from Proposition 22 that

$$\frac{\sum_{a_i \in A_i} \left( \left[ r_{i,a_i}(x) \right]^+ \right)^2}{1 + \sum_{a_i \in A_i} \left[ r_{i,a_i}(x) \right]^+} = u_i \left( \varphi_i(x), x_{-i} \right) - u_i(x) = u_i \left( x_i, x_{-i} \right) - u_i(x) = 0.$$
 (13)

Hence for all  $i \in \mathcal{I}$ ,  $a_i \in \mathcal{A}_i$ ,  $[r_{i,a_i}(x)]^+ = 0$  and hence  $u_i(a_i, x_{-i}) - u_i(x) \le 0$ . As a consequence, the strategy x is a NE.

Using Brouwer's fixed-point theorem (with the continuous map  $\phi$  defined over the nonempty convex compact set  $\prod_{i=1}^N \Delta(\mathcal{A}_i)$ ) together with Proposition 23 concludes the proof of Nash's theorem, i.e. guarantees the existence of NE.

# 5 Next lecture

In the next lecture, we will focus on how to learn NE under repeated interactions in the online learning setting for the fundamental game class of potential games.

### References

- Ky Fan. Fixed-point and minimax theorems in locally convex topological linear spaces. *Proceedings of the National Academy of Sciences*, 38(2):121–126, 1952.
- Gabriele Farina. Topics in multi-agent learning (mit 6.s890), lecture 2. https://www.mit.edu/~gfarina/2024/65890f24\_L02\_nfg\_nash/, 2024.
- Irving L Glicksberg. A further generalization of the kakutani fixed point theorem, with application to nash equilibrium points. *Proceedings of the American Mathematical Society*, 3(1): 170–174, 1952.
- Shizuo Kakutani. A generalization of Brouwer's fixed point theorem. *Duke Mathematical Journal*, 8(3):457 459, 1941. doi: 10.1215/S0012-7094-41-00838-4. URL https://doi.org/10.1215/S0012-7094-41-00838-4.
- Rida Laraki, Jérôme Renault, and Sylvain Sorin. *Mathematical foundations of game theory*, volume 202. Springer, 2019.
- Jason R. Marden. Game theory and multiagent systems (ece 149), lectures 11, 12, 14. https://web.ece.ucsb.edu/~jrmarden/ece149.html, 2020.
- Panayotis Mertikopoulos. *Online optimization and learning in games: Theory and applications*. PhD thesis, Grenoble 1 UGA-Université Grenoble Alpes, 2019.
- John F. Nash. Equilibrium points in *n*-person games. *Proceedings of the National Academy of Sciences*, 36(1):48–49, 1950. doi: 10.1073/pnas.36.1.48. URL https://www.pnas.org/doi/abs/10.1073/pnas.36.1.48.
- John F. Nash. Non-cooperative games. *Annals of Mathematics*, 54(2):286–295, 1951. URL http://www.jstor.org/stable/1969529.